

# A tiny defect, a terrible disease, a new approach

News from  
**The American Society  
for Cell Biology**  
**51st Annual Meeting**  
Denver, Colorado  
December 3–7, 2011

**EMBARGOED  
FOR RELEASE**  
10:00 am, U.S. Mountain Time  
Tuesday, December 6, 2011

## Contact

**Heather Ward, PhD**  
Department of Pathology  
University of New Mexico  
MSC08-4640  
225 Camino de Salud, CRF 225  
Albuquerque, NM 87131-5301  
505-272-1460  
hhward@unm.edu

## Author presents

Tuesday, December 6, 2011  
12:30 pm–2:00 pm  
Session: Post-Golgi Trafficking  
Presentation: 1628  
Board Number: B475  
Exhibit Halls: A/B/E/F

*Relaxin Improves Cystic  
Kidney Disease in Rats and  
Alters Transcription of  
Ciliary Trafficking Genes*

**H. Ward,<sup>1</sup> L. Danielson,<sup>1</sup> G.  
Pickett,<sup>2</sup> S. Ness,<sup>2</sup> C. Phillips,<sup>3</sup>  
B. Cowley,<sup>4</sup> A. Wandering-  
Ness<sup>1</sup>**


- 1 Department of Pathology, University of New Mexico, Albuquerque, NM
- 2 Department of Molecular Genetics and Microbiology, University of New Mexico, Albuquerque, NM
- 3 Department of Pathology and Laboratory Medicine, Indiana University School of Medicine, Indianapolis, IN
- 4 College of Medicine, Oklahoma University Health Sciences Center, Oklahoma City, OK

## A hormone that dilates blood vessels during pregnancy opens an indirect pathway for treating polycystic kidney disease

For something so seemingly inconsequential, primary cilia seemed an unlikely cause for something so devastating. Primary cilia are microscopic hairlike structures that stick out like antennae from virtually every cell in the human body, including the epithelial cells lining the kidneys. These nonmoving primary cilia were thought to be little more than evolutionary leftovers until 2000, when research first linked cilia defects to polycystic kidney disease (PKD), a life-threatening genetic disorder that affects 600,000 Americans, according to the National Institute of Diabetes, Digestive, and Kidney Diseases (NIDDK) and the Polycystic Kidney Disease Foundation. Half of those diagnosed with PKD will progress to end-stage renal disease by age 60.

A decade after the discovery, we know a tremendous amount about ciliopathies, a raft of diseases or conditions now associated with defects in primary cilia. Yet there is still no effective treatment for PKD. Researchers have concentrated on halting or reversing the formation of the characteristic large fluid-filled cysts that gradually encroach on kidney function, but Heather Ward, Angela Wandering-Ness, and others at the University of New Mexico, plus collaborators elsewhere, took another approach. They examined the noncystic aspects of PKD progression, particularly the poor blood flow and extensive internal scarring, called fibrosis, that encroaches on the glomeruli, the vital clusters of looping blood vessels that filter wastes and excess water from the blood. The group looked at a hormone called relaxin, first identified in pregnant women (but also present in men) that works (among other things) as a powerful vasodilator. What would be relaxin's effect on a cystic kidney?


Ward turned to special lab rats genetically altered to model PKD and saw an



The kidneys of special lab rats genetically altered to model PKD show the characteristic large fluid-filled cysts that gradually encroach on kidney function.

immediate improvement in renal function and blood flow after a course of relaxin. During human pregnancies, the relaxin hormone also regulates nitric oxide production, controls several signaling pathways, and stimulates the degradation of extracellular matrix around organ tissues by dissolving collagen. After four weeks on relaxin, the PKD model rats had lower collagen scores in their kidneys, indications that either scar formation was slowing down or the old fibroid tissue was breaking up. Most surprisingly, cysts seemed smaller in treated animals.

That finding sent Ward and colleagues exploring the differences in kidney gene expression between relaxin- and control-treated rats. The researchers now believe that relaxin, in part, affects genes associated with epithelial trafficking. Their hypothesis is that relaxin's direct effect on signaling pathways of kidney fibroblasts and vascular cells is improving the renal environment, indirectly affecting cystic epithelia and slowing cyst growth.

In any case, Ward and colleagues say that the dramatic improvement in renal function and decreased fibrosis they have seen in the rat models make relaxin a promising candidate for treating PKD, the first ciliopathy and a condition that still exacts a heavy toll in health costs and in human lives. 

*This work is funded by the National Institutes of Health (NIH)/National Center for Research Resources 5P20RR016480-10 UNM (PI, Griffith; Fellow, Ward), IDEA Networks of Biomedical Research Excellence (INBRE) Scholarship, and NIH/NIDDK R01 DK50141-11A1 (PI, Wandering-Ness), "Pathologic Mechanisms of Polycystic Kidney Disease." Ward is also supported by award number K12GM088021 from the National Institute of General Medical Sciences (PI, Wandering-Ness).*